



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

backwards to the middle of the costal edge and another to the middle of the dorsal edge; a third broad branch, connected with the central blotch only at one corner goes perpendicularly down to dorsal edge just before termen and a fourth equally narrowly connected branch reaches costal edge at apical fourth and sends a spur downwards and outwards, which nearly reaches the terminal edge. Just before apex are a few transverse black striæ. It might be more convenient to describe the duller black speckled color as ground color on which then the golden saffron color shows as there costal and there dorsal finger-like recurved spots nearly, but not quite meeting in the middle of the wing. Hind wings light golden saffron. On the under side of the forewings the dark design is slightly indicated and emphasized on costal edge by dark striation; hind wings have on the under side some scattered costal and apical striation. Abdomen and legs dull saffron. Alar expanse, female, 30-33 mm.

Venation typical with 12 veins in forewings; 7 and 8 separate; 7 to termen; 2 from before apical fourth of cell. Hind wings with 3 and 4 connate, 6 and 7 closely approximate; submedian vein not hairy.

*Habitat.* — Orizaba and Jalapa, Mexico (Wm. Schaus collection).

*Type.* — U. S. Nat. Mus., No. 10926.

***Tortrix auricomana*, new species.**

Labial palpi and tongue deep golden saffron; antennæ golden yellow, annulated with fuscous. Face, head and thorax deep golden yellow. Fore wings light straw yellow, evenly striated with thin black transverse lines and with entire costal, terminal and dorsal edge golden saffron; reaching from base of wing to apical third is a golden yellow streak along the subcostal vein and through the middle of the wing is a similar longitudinal streak beginning just before the end of the cell and terminating shortly before terminal edge of the wing; below this streak is a round golden saffron dot. Cilia golden. Hind wings light saffron yellow with a deep yellow hair pencil on vein 1b; submedian vein not hairy. Under side of both wings saffron yellow with costal and terminal parts slightly irrorate by short black transverse striæ. Abdomen and legs deep saffron yellow. Alar expanse, female, 30-34 mm.; male, 26-27 mm.

Venation as in the foregoing species.

*Habitat.* — Mexico City, Mexico. May and June (R. Müller, collector).

*Types.* — Male and female, U. S. National Museum, No. 10927.

---

A NOTE ON SYNONYMY.

Through the courtesy of Mr. A. Cosens, of Toronto, Canada, the U. S. National Museum has lately obtained topotypes of *Retinia austriana* Cosens (Can. Ent., XXXVIII, 362, 1906) in the author's own determination. The species proves to be a Phycitid, which Dr. H. G. Dyar has identified as *Pinipestis zimmermanni* Grote.—AUGUST BUSCK.